


# Max-N

Moxifloxacin 5 mg/ml · Ophthalmic Drops · 5 ml

**Main uses**

- Bacterial conjunctivitis (infection of the outer eye) caused by susceptible bacteria
- Eye infections in adults and children where an antibiotic eye drop is prescribed

PRODUCT INFORMATION

# About Max-N

Max-N contains moxifloxacin, a fluoroquinolone antibiotic. It stops bacteria from multiplying and so clears the infection.

Max-N eye drops are used to treat bacterial conjunctivitis (infection of the outer layer of the eye) caused by bacteria that are sensitive to moxifloxacin.

Max-N is a sterile eye drop for use in the eye only, as advised by a doctor.

# How to use

Use Max-N exactly as your doctor has told you, and finish the course even if the eye looks better.

1. The usual dose is one drop in the affected eye three times a day, for the number of days your doctor advises.
2. Wash your hands first. Tilt your head back, pull the lower eyelid down gently and put the drop in the pocket formed, then close the eye for a moment.
3. If you use other eye drops or ointments, wait at least 5 minutes between each medicine. Do not let the tip of the bottle touch the eye or anything else.
4. Do not wear contact lenses while you have an eye infection.

Do not swallow or inject Max-N. Throw the bottle away 28 days after first opening it.

# Composition

EYE DROPS

Each ml contains moxifloxacin 5 mg (as moxifloxacin hydrochloride IP). Sterile aqueous vehicle.

PACK SIZES

Bottle of 5 ml.

# Side effects and warnings

Like all medicines, Max-N can cause side effects, although not everybody gets them. The most common are eye irritation, eye pain, redness and a change in taste.

- Do not use Max-N if you are allergic to moxifloxacin, other quinolone antibiotics or any of the other ingredients.
- Stop using Max-N and tell your doctor straight away if you get a rash, swelling of the face or throat, or difficulty breathing.
- Long or repeated use can allow bacteria or fungi that are not sensitive to grow. Tell your doctor if the infection does not improve in a few days.
- Your vision may be blurred for a short while after the drop. Do not drive or use machines until it clears.
- If you are pregnant or breast-feeding, ask your doctor before using Max-N.

Always read the patient information leaflet in the pack. If you have any questions, ask your doctor or pharmacist.